


Canine Atrial KRCCN02-SM

Simucore Model Based Upon: Kneller, Ramirez, Chartier, Courtemanche, Nattel Model of Canine Atrial Cardiac Action Potentials, 2002; v. 1.4

Table of contents

1 Description.....	2
2 References.....	3
3 Ordering.....	3


1. Description



Example action potential after 10 iterations (BCL = 400 ms).

This model improves long-term stability of Ramirez, Nattel, Courtemanche Canine model by adding background Cl^- current and $\text{Na}^+\text{-Cl}^-$ cotransporter formulations. Maximal conductances of I_{Ca} , I_{L} , I_{to} and I_{ur} were also adjusted to stabilize action potential configurations during long runs.

The example action potential below was recorded after 50,000 model pacings at BCL=400 ms. The action potential configuration after this >5.5h long run is very similar to the action potential recorded after only 10 iterations (above). This model is therefore suitable for simulations where long-term ionic concentration stability is required over prolonged periods of time.



Example action potential after 50,000 iterations (BCL = 400 ms).

Abstract excerpt: *"The present study evaluated transients during long-term simulated activity in a mathematical model of the canine atrial AP."*

2. References

- Kneller J, Ramirez RJ, Chartier D, Courtemanche M, Nattel S. Time-dependent transients in an ionically based mathematical model of the canine atrial action potential. *Am J Physiol Heart Circ Physiol.* 2002 Apr;282(4):H1437-51. PMID: [11893581](https://pubmed.ncbi.nlm.nih.gov/11893581/)

3. Ordering

- [Order this model](#) or [request further information](#).